


38"

1.25"
SEAL

10 YEAR
GUARANTEE
WHEN APPLIED
AS DIRECTED



St. Gabriel
ORGANICS
SINCE 1973



ONLY 4 APPLICATIONS:
Spring, Summer & Fall for 2 years

YEAR 1: SPRING SUMMER FALL YEAR 2: SPRING SUMMER FALL

SPREADER SETTINGS

Use of this material works best when dispersed on lawn areas using a Drop Spreader.

Earthway EndSpreader 2500a Drop.....	5
Greenview Drop Spread	5
Republic Drop es 530	6
Scotts 1000 Aqueous Drop	5
Sears Drop	7
TruGreen Drop	6
Tu-Temper Drop	6
Ortho Drop	6
Amae Drop Spreader	2
37A Green Drop Spreader	2

All spreader settings are for Drop type spreaders. If your spreader is not listed, or, if your existing spreader is old and worn, we suggest checking the settings and calibrations by using a small amount of material.

ONE POUND OF MATERIAL WILL COVER 350 SQUARE FEET.

INOCULATION BY DROP SPREADER:

For lawns and other grassy areas around homes, spread MILKY SPORE GRANULAR evenly over the treatment area with a drop spreader at the rate of 1 lb per 350 sq ft or 20 lb per 7,000 sq ft. To achieve maximum build up of spores in the soil and best grub control, apply this product during the Spring, Summer and Fall for two successive years.

The additional applications are necessary to assure that the concentration of Milky Spore reaches levels high enough to initiate and maintain infection. Milky Spore infects and controls Japanese beetle grubs in the soil and, once established, may remain viable for many years.

Follow these application directions and use the specified drop spreader setting:

- Fill the drop spreader with MILKY SPORE GRANULAR.
- Apply uniformly over the treatment area with the spreader settings specified in the table below.
- Water the treated turf lightly after application to soak spores into the soil.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:
CAUTION: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID:

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Hot-Line Number 800-801-0061

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information or questions, call 1-800-801-0061, Monday through Friday 9am to 5pm Eastern time. During other times call your poison control center at 1-800-222-1222.

GRUB CONTROL
Milky Spore
GRANULAR


Milky Spore GRANULAR GRUB CONTROL



Adult
Beetle



Healthy
Grub



Magnified
Spores



Grub infected with
Milky Spore

READY TO USE

Rid your lawn of unsightly brown spots that may be caused by grub damage. Control Japanese Beetle grubs in the soil. Use on grass, in gardens and mulch beds.

To achieve best grub control, apply Milky Spore Spring, Summer and Fall for two years to achieve maximum build up of spores in the soil.

Milky spore can be applied any time the ground is not frozen. Once established, milky spore may remain viable for many years.

10 YEAR
GUARANTEE
WHEN APPLIED
AS DIRECTED

TREATS 7,000 SQUARE FEET

NET WEIGHT 20 LBS (9.12 KG)

CAUTION:

Keep Out of Reach of Children. See Precautionary Statements on back panel.

EPA Est. no. 63191-VA-1
EPA Reg. no. 63191-16

Milky Spore • GRANULAR • GRUB CONTROL

QUICK FACTS

KILLS: Japanese Beetle Grubs
COVERAGE: This bag covers up to 7,000 sq. ft.
WHERE TO USE: Lawns, Flower Beds, Ground Covers, Vegetable Gardens, and around Trees & Shrubs.
WHEN TO USE: Milky Spore Granular is a 2-year program. Apply Spring, Summer, & Fall for 2 consecutive years.
HOW MUCH TO USE: See spreader settings.
HOW TO REACH US: For questions or comments, call toll-free (800) 801-0061 or visit us on-line at: www.stgabrielorganics.com

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE ON LAWNS AND OTHER GRASSY AREAS AROUND HOMES, ORNAMENTAL GARDENS AND FLOWER BEDS, MULCHED BEDS AND ORNAMENTAL GROUND COVERS.

To control Japanese beetle grubs in lawns, turf or other areas specified on this label, apply MILKY SPORE GRANULAR during any season of the year when the ground is not frozen and water lightly after application to soak spores into the soil.

GRUBS:
• Apply Spring, Summer, & Fall for 2 consecutive years.
LAWNS & GARDENS, TREES & SHRUBS:
• How Much: See Spreader Settings.
• Coverage: To be sure you cover the entire area uniformly first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules.
• Water thoroughly after applying.

FOR BEST RESULTS:
• Apply early before grubs hatch.
• Water thoroughly within 24 hours after applying.
• Do not mow until the granules are worked into the soil.

SIGNS OF GRUB DAMAGE TO LAWNS:
Japanese Beetle grubs cause extensive damage to lawns by feeding on grass roots. Grass turns yellow, then brown, and dies. Grubs chew off roots so that the grass is easily pulled up in pieces in your hands. You can roll it back like a carpet to reveal the grubs. If patches of dead grass appeared in your lawn late summer, check for grubs.

Environmental Hazards:
To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when heavy rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto treated area of lawn or garden, will help prevent run off to water bodies or drainage systems.

STORAGE DISPOSAL
Do not contaminate water, food or feed by storage or disposal.
Keep container tightly closed to protect product from moisture.

PESTICIDE STORAGE Store in a cool, dry place away from direct sunlight.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:
Nonrefillable container. Do not reuse or refill container.
If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY:
To the extent consistent with applicable law, St. Gabriel Organics makes no other express or implied warranties, either of merchantability or fitness, for a particular use and shall not be liable for misuse or conditions beyond its control, such as handling or storage.

St. Gabriel Organics
14041 Littlefield Dr., Orange, VA 22660
1-800-801-0061
www.StGabrielOrganics.com