



PYRECAN™ INSECTICIDE

READY TO USE

**KILLS MITES, WHITEFLIES, APHIDS, AND ALL
ACCESSIBLE STAGES OF LISTED INSECTS & EGGS**

KILLS INSECTS ON CONTACT

MAY BE USED UP TO AND INCLUDING DAY OF HARVEST



Keep Out Of Reach Of Children

CAUTION

(See booklet for Additional Precautionary
Statements and Directions for Use)

Net Contents 128 FL. OZ. (3.785 L.)

XXXXXXXX

ACTIVE INGREDIENTS:

Pyrethrins 0.01%

Canola Oil 1.00%

OTHER INGREDIENTS: 98.99%

TOTAL: 100.00%

This product contains 0.0824 lbs. of canola oil and
0.0008 lbs. of pyrethrins per gallon.

EPA Est. No. 4-NY-1 EPA Reg. No. 67702-35-93133

99000

PYRECAN™ INSECTICIDE READY TO USE

KILLS ALL STAGES OF INSECT GROWTH-EGG TO ADULT

FOR USE ON: VEGETABLES, FRUITS, ORNAMENTALS,
SHRUBS, TREES AND HOUSEPLANTS

KILLS 100+ LISTED INSECTS ON CONTACT

FOR INDOOR AND OUTDOOR USES

Manufactured under a license of
W. Neudorff GmbH KG, Germany.
EPA Reg. No. 67702-35-93133 EPA Est. No. 4-NY-1

Distributed by: A.M. Leonard
241 Fox Drive, Piqua, OH 45356
Gallon



8 5000145016 4

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label. Use only as directed. Shake well before each use.

Use Restrictions

- Do not make any applications during rain.
- Do not water the treated area to the point of run-off.
- Do not wet plants to point of run-off or drip.
- Do not allow adults, children or pets to enter the treated area until sprays have dried.
- Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.
- Remove pets, birds, and cover fish aquariums before spraying.
- Do not apply within 21 days of a sulfur application.
- Do not use this product when temperatures are above 90°F.
- Do not apply more than 10 times per season.
- Do not reapply within 3 days except under extreme pest pressure.
- Do not apply more than 1 time per day.
- In case of extreme pest pressure do not reapply within 24 hours.
- Do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during application.

Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.

Other than applications to building foundations, all applications to impervious surfaces, such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

To prevent product run-off, do not over water treated areas or apply prior to heavy rainfall.

Phytotoxicity Note: Oil removes the glaucous (blue) bloom from such evergreens as Colorado blue spruce and Koster spruce. Apply carefully to blooms and buds and on oil sensitive plants such as: Chamaecyparis, Cryptomeria, Japanese Holly, Red Cedar and Smoke Tree. If concerned about sensitivity of plants, test by spraying a portion of the plant and wait several days to determine if burn occurs before spraying larger areas.

APPLICATION INSTRUCTIONS

Thoroughly wet all parts of the plants. Do not wet plants to point of runoff or drip. Repeat treatment in 1-2 weeks if needed. Do not apply more than 20 fl. oz. per 100 square feet per application. May be used on edible crops up to and including day of harvest.

PESTS/SITES

This product kills exposed: Adelgids, Ants including Foraging Ants, Sweet-Feeding Ants, Pavement Ants, Moisture Ants, Little Black Ants, Argentine Ants, Crazy Ants, Acrobat Ants, Bigheaded Ants, Cornfield Ants, Ghost Ants, Odorous House Ants, Thief Ants, Texas Leafcutting Ants, Field Ants, and other Common Ants (except Fire Ants, Harvester Ants, Carpenter Ants, and Pharaoh Ants), Aphids (including Green Peach Aphids), Armyworms (including Beet Army Worms, Western Yellow Striped Armyworms), Bagworms, Beetles (Colorado potato beetle, flea beetle, Japanese beetle, asparagus beetle, Blister Beetles, 12-spotted Cucumber Beetles, Cucumber Beetle, Darkling Beetles (lesser mealworm), Saw-toothed Grain Beetles, Elm Leaf Beetles, Bean Beetles, Mexican Bean Beetles), Borers (including European Corn Borer, Branch and Twig Borers, Shot hole borers and Pacific flathead borers), Box Elder Bugs, Cabbageworms (including Cross-striped Cabbageworm and Imported Cabbageworm), Cankerworms, Caterpillars (Gypsy Moth Caterpillars, Tent Caterpillar, Eastern Tent Caterpillar, Forest Tent Caterpillar and Tomato Fruitworm Caterpillar), Corn Earworms, Crane Flies, Crickets, Diamondback Moth Larvae, Earwigs, Fireworms, Fungus Gnats, Fruit Flies (Including Cherry Fruit Fly), Glassy Winged Sharpshooter, Grape Leaf Skeletonizer, Grasshoppers,

Green Fruit Worm, Harlequin Bugs, Hornworms (including Tomato Hornworm), Katydid, Lace Bugs, Leafhoppers (including Grape Leafhopper, and Potato Leafhoppers), Leafminers, Leafrollers (including Fruittree Leafrollers), Leaf-tiers, Loopers (including Cabbage Looper), Lygus, Maggots (including Apple Maggot, Cabbage Maggots, and Onion Maggot), Mealybugs (including Citrus Mealybugs, Grape Mealybugs and Vine Mealybugs), Midges, Millipedes, Mites (including Clover Mites, European Red Mite and Spider Mites), Moths (including Artichoke Plume Moth, Codling Moth, Diamondback Moth, European Pine Tip Moth, Gypsy Moths, Indian Meal Moth, Mediterranean Flour Moth), Mushroom Flies, Navel Orangeworm, Plant Bugs (including Leaf-footed Plant Bug, Tarnished Plant Bug), Psyllids (including Pear Psyllids), Sawfly Larvae (pear and rose slugs), Scale insects (including Soft Scale and Armoured Scale), Silverfish, Skippers, Sowbugs, Spittlebugs, Squash Bugs, Stink Bugs, Thrips (including Greenhouse Thrips), Vinegar Flies, Webworms, Weevils (including Boll Weevil, Carrot Weevil, Clover Weevil, Rice Weevil, Pepper Weevil), Whitefly and Phylloxera.

VEGETABLES AND HERBS: Use as directed above on vegetables and herbs such as asparagus, beans, broccoli, Brussels sprouts, cabbage, cauliflower, kale, oriental vegetables,

Herbs, lettuce, eggplant, peppers, tomatoes, potatoes, cucumbers, melons and squash.

FRUITS AND NUTS: Use as directed above on fruits and nuts such as apples, avocados, citrus (grapefruit, lemons, oranges, tangerines, limes), peaches, nectarines, apricots, pears, almonds, pecans, walnuts, kiwi fruit, blackberries, loganberries, raspberries, strawberries and grapes.

ORNAMENTAL TREES AND SHRUBS: Use as directed above on ornamental trees and shrubs such as azaleas, birch, camellias, dogwood, holly, rhododendrons, evergreens and shade trees. See PHYTOTOXICITY section for additional use restrictions.

FLOWERS: Use as directed above on flowers such as asters, carnations, chrysanthemums, geraniums, marigolds, petunias and roses.

HOME GREENHOUSE USE: Use as directed above in home greenhouses on flowering and foliage plants; vegetables and fruits.

INDOOR USE: For use indoors on houseplants as directed above.

DORMANT USE: Kills exposed egg stages of pests present in dormant season and pests on dormant shrubs (roses), evergreens, fruit trees and shade trees. Thoroughly wet the dormant plants or trees.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING & DISPOSAL: Nonrefillable. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run-off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the area.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the poison control center 1-800-222-1222.

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.